



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

of the aorta. Having lately had the opportunity of dissecting another specimen which died in the Zoological Gardens of this city, he called attention to a slightly different disposition of these vessels. In this case an innominate gave off the right subclavian, right vertebral, the left subclavian, and left vertebral separately, and then terminated in the two common carotids. This arrangement also differs from that described by Prof. Owen, thus illustrating the variability in the origin of the bloodvessels in these animals. The recurrent laryngeal nerve was found as described by the eminent anatomist just mentioned, that is, the lower two-thirds almost completely atrophied, while the upper third came from the pneumogastric just below the superior laryngeal.

On the Habits of Quiscalus purpureus.—JOSEPH WILLCOX spoke of the custom of some crow blackbirds (*Quiscalus purpureus*) of eating fish, and their method of catching them through the intervention of another animal.

Last winter when on the shore of the St. John's River, near Lake Jessup, in Florida, he observed an unusual commotion among the crow blackbirds, which were congregating near the water, and anxiously looking into it. Soon a loud splash and noise were made in the vicinity by a large bass in making a charge upon his favorite food, the small fry, which, in their frantic efforts to escape, jumped out of the water, and many of them fell upon the land.

The blackbirds, which were evidently experts at the game, immediately pounced upon the small fish and swallowed them before they had time to hop back into the water. This performance was repeated many times.

On one occasion he examined the stomach of a bass in order to ascertain what it had taken for dinner. In it he found a rat that had evidently been imprisoned only a short time.

FEBRUARY 27.

The President, Dr. RUSCHENBERGER, in the chair.

Forty-two members present.

Frank L. Scribner and Mrs. Mary Wagner were elected members.

Prof. Alphonse Favre of Geneva, Prof. Paul De Loriol of Geneva, Baron Ferdinand Freiherrn Von Richthofen of Berlin, Dr. Ferd. Von Hochstetter of Vienna, Prof. Edouard Desor of Neuchâtel, Prof. Ludwig Rüttimeyer of Basel, Prof. Valerien De Mueller of St. Petersburg, H. Trautschold of Moscow, and Lieut. A. W. Vodges, U. S. A., were elected correspondents.

The following papers were ordered to be printed:—